

ALGEBRAIC CYCLES ON SEVERI-BRAUER SCHEMES OF PRIME DEGREE OVER A CURVE

CRISTIAN D. GONZÁLEZ-AVILÉS

ABSTRACT. Let k be a perfect field and let p be a prime number different from the characteristic of k . Let C be a smooth, projective and geometrically integral k -curve and let X be a Severi-Brauer C -scheme of relative dimension $p - 1$. In this paper we show that $CH^d(X)_{\text{tors}}$ contains a subgroup isomorphic to $CH_0(X/C)$ for every d in the range $2 \leq d \leq p$. We deduce that, if k is a number field, then $CH^d(X)$ is finitely generated for every d in the indicated range.

1. INTRODUCTION.

Let k be a perfect field with algebraic closure $\bar{k}$. Very little is known about algebraic cycles on algebraic k -varieties, especially in codimension greater than 2 or dimension greater than zero. Let p be a prime number different from the characteristic of k and let C be a smooth, projective and geometrically integral k -curve. In this paper we study a certain subgroup of $CH^d(X)_{\text{tors}}$ for a Severi-Brauer C -scheme $q: X \rightarrow C$ of relative dimension $p - 1$ and any integer d such that $2 \leq d \leq p$. Let

$$CH_0(X/C) = \text{Ker} \left[CH_0(X) \xrightarrow{q_*} CH_0(C) \right]$$

and let $\pi^*: CH^d(X) \rightarrow CH^d(\bar{X})$ be induced by the extension-of-scalars map $\bar{X} \rightarrow X$, where $\bar{X} = X \otimes_k \bar{k}$. Then the following holds.

Main Theorem. *For any d as above, there exists a canonical isomorphism*

$$\text{Ker} \left[CH^d(X) \xrightarrow{\pi^*} CH^d(\bar{X}) \right] \simeq CH_0(X/C).$$

Consequently, if k is a number field, then $CH^d(X)$ is finitely generated.

2000 *Mathematics Subject Classification.* Primary 14C25; Secondary 14C15 .

Key words and phrases. Algebraic cycles, Chow groups, curves, Severi-Brauer schemes.

The author is partially supported by Fondecyt grant 1061209 and Universidad Andrés Bello grant DI-29-05/R.

ACKNOWLEDGEMENT.

I thank B.Kahn for some helpful comments.

2. PRELIMINARIES.

Let k be a perfect field, fix an algebraic closure $\bar{k}$ of k and let $\Gamma = \text{Gal}(\bar{k}/k)$. Now let C be a smooth, projective and geometrically integral k -curve and let X be a Severi-Brauer scheme over C [4, §8]. There exists a proper and flat k -morphism $q: X \rightarrow C$ all of whose fibers are Severi-Brauer varieties of dimension $m - 1$ ($m \geq 1$) over the appropriate residue field [loc.cit.]. We will write X_η for the generic fiber $X \times_C \text{Spec } k(C)$ of q and A for the central simple $k(C)$ -algebra associated to X_η . We define

$$CH_0(X/C) = \text{Ker} \left[CH_0(X) \xrightarrow{q_*} CH_0(C) \right].$$

Now let C_0 be the set of closed points of C . The group of *divisorial norms* of X/C (cf. [6]) is the group

$$k(C)_{\text{dn}}^* = \{f \in k(C)^*: \forall y \in C_0, \text{ord}_y(f) \in (q_y)_*(CH_0(X_y))\}$$

where, for each $y \in C_0$, $q_y: X_y \rightarrow \text{Spec } k(y)$ is the structural morphism of the fiber X_y . This group is closely related to $CH_0(X/C)$ (see [2, Proposition 3.1]). Indeed, there exists a canonical isomorphism

$$CH_0(X/C) \simeq k(C)_{\text{dn}}^* / k^* \text{Nrd } A^*.$$

Now fix an integer d such that $1 \leq d \leq m$ and let

$$CH^d(X)' = \text{Ker} \left[CH^d(X) \xrightarrow{\pi^*} CH^d(\bar{X})^\Gamma \right],$$

where $\pi: \bar{X} \rightarrow X$ is the canonical map. A simple transfer argument shows that $CH^d(X)'$ is a subgroup of $CH^d(X)_{\text{tors}}$. Now, since $\bar{X} \rightarrow \bar{C}$ has a section, $\bar{X}$ is a projective bundle over $\bar{C}$. Thus there exists an isomorphism

$$CH^d(\bar{X}) \simeq \mathbb{Z} \oplus CH_0(\bar{C}).$$

(see [3, Theorem 3.3(b), p.34]). Therefore, if $J_C(k)$ is finitely generated, where J_C is the Jacobian variety of C (e.g., k is a number field or $C = \mathbb{P}_k^1$), then $CH^d(X)$ is finitely generated if and only if $CH^d(X)'$ is finite.

3. THE GENERAL METHOD.

Let C be as above and let X be any smooth, projective and geometrically integral k -variety such that there exists a proper and flat morphism $q: X \rightarrow C$ whose generic fiber X_η is geometrically integral. We have an exact sequence [7]

$$(1) \quad H^{d-1}(X_\eta, \mathcal{K}_d) \xrightarrow{\delta} \bigoplus_{y \in C_0} CH^{d-1}(X_y) \rightarrow CH^d(X) \xrightarrow{j^*} CH^d(X_\eta) \rightarrow 0,$$

where $j: X_\eta \rightarrow X$ is the natural map and the map which we have labeled δ will play a role later when $k = \bar{k}$. A similar exact sequence exists over $\bar{k}$, and we have two natural exact commutative diagrams:

$$\begin{array}{ccccccc} 0 & \longrightarrow & \text{Ker } j^* & \longrightarrow & CH^d(X) & \longrightarrow & CH^d(X_\eta) \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & (\text{Ker } \bar{j}^*)^\Gamma & \longrightarrow & CH^d(\bar{X})^\Gamma & \longrightarrow & CH^d(\bar{X}_\eta)^\Gamma \end{array}$$

and

$$(2) \quad \begin{array}{ccccccc} 0 & \longrightarrow & \frac{H^{d-1}(X_\eta, \mathcal{K}_d)}{j^* H^{d-1}(X, \mathcal{K}_d)} & \longrightarrow & \bigoplus_{y \in C_0} CH^{d-1}(X_y) & \longrightarrow & \text{Ker } j^* \\ & & \downarrow & & \downarrow \oplus_{\bar{y}|y} \pi_{\bar{y}}^* & & \downarrow \\ 0 & \longrightarrow & \left(\frac{H^{d-1}(\bar{X}_\eta, \mathcal{K}_d)}{\bar{j}^* H^{d-1}(\bar{X}, \mathcal{K}_d)} \right)^\Gamma & \xrightarrow{\bar{\delta}} & \bigoplus_{\bar{y}|y} CH^{d-1}(\bar{X}_{\bar{y}})^{\Gamma_y} & \longrightarrow & (\text{Ker } \bar{j}^*)^\Gamma \end{array}$$

where, for each $y \in C_0$, we have fixed a closed point $\bar{y}$ of $\bar{C}$ lying above y and written $\Gamma_y = \text{Gal}(\bar{k}/k(y))$. Set

$$CH^d(X_\eta)' = \text{Ker} \left[CH^d(X_\eta) \rightarrow CH^d(\bar{X}_\eta)^\Gamma \right]$$

and, for each $y \in C_0$,

$$CH^{d-1}(X_y)' = \text{Ker} \left[CH^{d-1}(X_y) \xrightarrow{\pi_{\bar{y}}^*} CH^{d-1}(\bar{X}_{\bar{y}})^{\Gamma_y} \right].$$

Now define

$$(3) \quad E(\bar{X}/\bar{C}) = \text{Coker} \left[\frac{H^{d-1}(X_\eta, \mathcal{K}_d)}{j^* H^{d-1}(X, \mathcal{K}_d)} \longrightarrow \left(\frac{H^{d-1}(\bar{X}_\eta, \mathcal{K}_d)}{\bar{j}^* H^{d-1}(\bar{X}, \mathcal{K}_d)} \right)^\Gamma \right].$$

Then, applying the snake lemma to the preceding diagrams, we obtain¹

Proposition 3.1. *There exists a natural exact sequence*

$$\begin{aligned} \bigoplus_{y \in C_0} CH^{d-1}(X_y)' &\rightarrow \text{Ker} [CH^d(X)' \rightarrow CH^d(X_\eta)'] \\ &\rightarrow \text{Ker} \left[E(\overline{X}/\overline{C}) \rightarrow \bigoplus_{y \in C_0} \frac{CH^{d-1}(\overline{X}_{\overline{y}})^{\Gamma_y}}{\pi_{\overline{y}}^* CH^{d-1}(X_y)} \right] \rightarrow 0, \end{aligned}$$

where $E(\overline{X}/\overline{C})$ is the group (3).

As regards the right-hand group in the exact sequence of the proposition, the following holds. Let

$$H^{d-1}(X_\eta, \mathcal{K}_d)' = \text{Im} \left[H^{d-1}(X_\eta, \mathcal{K}_d) \rightarrow H^{d-1}(\overline{X}_{\overline{\eta}}, \mathcal{K}_d)^\Gamma \right]$$

and

$$\text{Sal}_d(X/C) = \left\{ f \in H^{d-1}(\overline{X}_{\overline{\eta}}, \mathcal{K}_d)^\Gamma : \forall y \in C_0, \bar{\delta}_{\overline{y}}(f) \in \pi_{\overline{y}}^* CH^{d-1}(X_y) \right\},$$

where $\bar{\delta}$ and $\pi_{\overline{y}}^*$ are the maps of diagram (2).

Proposition 3.2. *There exists a natural exact sequence*

$$\begin{aligned} 0 &\rightarrow \frac{\text{Sal}_d(X/C)}{(\bar{j}^* H^{d-1}(\overline{X}, \mathcal{K}_d))^\Gamma \cdot H^{d-1}(X_\eta, \mathcal{K}_d)'} \\ &\rightarrow \text{Ker} \left[E(\overline{X}/\overline{C}) \rightarrow \bigoplus_{y \in C_0} \frac{CH^{d-1}(\overline{X}_{\overline{y}})^{\Gamma_y}}{\pi_{\overline{y}}^* CH^{d-1}(X_y)} \right] \\ &\rightarrow H^1(\Gamma, \bar{j}^* H^{d-1}(\overline{X}, \mathcal{K}_d)). \end{aligned}$$

Proof. This follows by applying the snake lemma to a diagram of the form

$$\begin{array}{ccccccc} 0 & \longrightarrow & A & \longrightarrow & B & \longrightarrow & B/A \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & \overline{A}^\Gamma & \longrightarrow & \overline{B}^\Gamma & \longrightarrow & (\overline{A}/\overline{B})^\Gamma \longrightarrow H^1(\Gamma, \overline{A}) \end{array}$$

with $\overline{A} = \bar{j}^* H^{d-1}(\overline{X}, \mathcal{K}_d)$, $\overline{B} = H^{d-1}(\overline{X}_{\overline{\eta}}, \mathcal{K}_d)$, etc. □

¹ Proposition 3.1 was inspired by [1, Proposition 1.1].

4. PROOF OF THE MAIN THEOREM.

Let C and A be as in Section 2, let p be a prime number different from the characteristic of k and let X be a Severi-Brauer scheme over C of relative dimension $p - 1$.

Lemma 4.1. *There exists a Γ -isomorphism*

$$\bar{j}^* H^{d-1}(\bar{X}, \mathcal{K}_d) \simeq \bar{k}^*.$$

Proof. Clearly, $\bar{j}^* H^{d-1}(\bar{X}, \mathcal{K}_d)$ is the kernel of the map

$$\bar{\delta}: H^{d-1}(\bar{X}_{\bar{\eta}}, \mathcal{K}_d) \rightarrow \bigoplus_{\bar{y}|y} CH^{d-1}(\bar{X}_{\bar{y}})$$

appearing in the exact sequence (1) over $\bar{k}$. Now $\bar{X}_{\bar{\eta}} \simeq \mathbb{P}_{\bar{\eta}}^{p-1}$ and $\bar{X}_{\bar{y}} \simeq \mathbb{P}_{\bar{k}}^{p-1}$ for every $\bar{y}$, whence we have Γ -isomorphisms

$$H^{d-1}(\bar{X}_{\bar{\eta}}, \mathcal{K}_d) \simeq \bar{k}(C)^*$$

and

$$CH^{d-1}(\bar{X}_{\bar{y}}) \simeq \mathbb{Z}$$

for each $\bar{y}$. Under these isomorphisms, the map $\bar{\delta}$ above corresponds to the canonical map

$$\begin{aligned} \bar{k}(C)^* &\rightarrow \bigoplus_{\bar{y}|y} \mathbb{Z}, \\ f &\mapsto (\text{ord}_{\bar{y}}(f))_{\bar{y}|y}, \end{aligned}$$

which yields the lemma. $\square$

Theorem 4.2. *For every d such that $2 \leq d \leq p$, there exists a canonical isomorphism*

$$CH^d(X)' \simeq CH_0(X/C).$$

Proof. By Lemma 4.1, Hilbert's Theorem 90 and Proposition 3.2, there exists a natural isomorphism

$$\text{Ker} \left[E(\bar{X}/\bar{C}) \rightarrow \bigoplus_{y \in C_0} \frac{CH^{d-1}(\bar{X}_{\bar{y}})^{\Gamma_y}}{\pi_{\bar{y}}^* CH^{d-1}(X_y)} \right] \simeq \frac{\text{Sal}_d(X/C)}{k^* H^{d-1}(X_{\eta}, \mathcal{K}_d)' }.$$

On the other hand, by [5, (8.7.2)], $H^{d-1}(X_{\eta}, \mathcal{K}_d)' = \text{Nrd } A^*$ for every d such that $2 \leq d \leq p$ and, for each $y \in C_0$,

$$\pi_{\bar{y}}^* CH^{d-1}(X_y) \simeq \pi_{\bar{y}}^* CH^{p-1}(X_y) \simeq (q_y)_* CH_0(X_y) \quad (= \mathbb{Z} \text{ or } p\mathbb{Z}).$$

The latter implies that $\text{Sal}_d(X/C) = k(C)_{\text{dn}}^*$, whence

$$\begin{aligned} \text{Sal}_d(X/C)/k^* H^{d-1}(X_{\eta}, \mathcal{K}_d)' &\simeq k(C)_{\text{dn}}^*/k^* \text{Nrd } A^* \\ &\simeq CH_0(X/C). \end{aligned}$$

Finally, [loc.cit.] shows that the groups $CH^d(X_\eta)$ and $CH^{d-1}(X_y)$ ($y \in C_0$) are torsion free, whence $CH^d(X_\eta)'$ and $CH^{d-1}(X_y)'$ vanish. The theorem now follows from Proposition 3.1. $\square$

Corollary 4.3. *Let d be such that $2 \leq d \leq p$. Then $CH^d(X)'$ is finite if*

- (1) *k is a number field, or*
- (2) *k is a field of finite type over $\mathbb{Q}$, $C = \mathbb{P}_k^1$ and X has a 0-cycle of degree one.*

In each of these cases, the group $CH^d(X)$ is finitely generated.

Proof. Indeed, in these cases the group $CH_0(X/C)$ is finite [2]. $\square$

REFERENCES

- [1] Colliot-Thélène, J.-L. and Skorobogatov, A.: *Groupe de Chow des zéro-cycles sur les fibrés en quadriques*. *K-Theory* **7**, pp. 477-500 (1993).
- [2] Frossard, E.: *Groupe de Chow de dimension zéro des fibrations en variétés de Severi-Brauer*. *Comp. Math.* **110**, pp. 187-213 (1998).
- [3] Fulton, W.: *Intersection Theory*. Second Ed. Springer-Verlag, 1998.
- [4] Grothendieck, A.: *Le Groupe de Brauer I*. In: *Dix Exposés sur la Cohomologie des Schémas*. North-Holland, Amsterdam, pp. 46-66 (1968).
- [5] Merkurjev, A.S. and Suslin, A.A.: *K-cohomology of Severi-Brauer varieties and the norm residue homomorphism*. *Math. USSR Izv.* **21**, No. 2, pp. 307-340 (1982).
- [6] Salberger, P.: *Galois descent and class groups of orders*. *Lect. Notes in Math.* **1142**, pp. 239-255 (1985).
- [7] Sherman, C.: *Some theorems on the K-Theory of coherent sheaves*. *Comm. in Alg.* **7**(14), pp. 1489-1508 (1979).

DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD ANDRÉS BELLO, CHILE
E-mail address: cristianguzalez@unab.cl